



Search:

[Email this page](#)
[Printer-friendly version](#)

View By Topics **Costs of Falls Among Older Adults**

A-Z Injury Topics

CDC Injury Fact Book

Data & Statistics

Fact Sheets

Funding and Research

Publications

Injury Response Overview

- [Acute Injury Care](#)
- [Alcohol Screening and Brief Intervention](#)
- [Mass Casualties State Programs](#)
- [Traumatic Brain Injury](#)

Violence Prevention Overview

- [Child Maltreatment](#)
- [Intimate Partner Violence](#)
- [Sexual Violence](#)
- [Suicide](#)
- [Youth Violence](#)

Unintentional Injury Prevention Overview

- [Child Passenger Safety](#)
- [Preventing Falls Among Older Adults](#)
- [Fire Deaths and Injuries](#)
- [Fireworks Injury Prevention](#)
- [Impaired Driving](#)
- [Older Adult Drivers](#)
- [Playground Injuries](#)
- [Teen Drivers](#)
- [Water-Related Injuries](#)

Information Resource

- [Calendar](#)
- [Related Web Sites](#)

About CDC Injury

How big is the problem?

- More than one third of adults 65 and older fall each year (Hornbrook et al. 1994; Hausdorff et al. 2001).
- Of those who fall, 20% to 30% suffer moderate to severe injuries that make it hard to get around or live alone and increase the chance of early death (Alexander et al. 1992).
- Older adults are hospitalized for fall-related injuries five times more often than they are for injuries from other causes (Alexander et al. 1992).



How do we calculate costs?

The costs of fall-related injuries are often shown in terms of direct costs.

- Direct costs are what patients and insurance companies pay for the treatment of fall-related injuries. These costs include fees for hospital and nursing home care, doctors and other professional services, rehabilitation, community-based services, use of medical equipment, prescription drugs, changes made to the home, and insurance processing (Englander et al. 1996).
- Direct costs do not account for the long-term effects of these injuries, such as disability, dependence on others, lost time from work and household duties, or reduced quality of life.

How costly are fall-related injuries among older adults?

- A study of people 72 and older found that the average health care cost of a fall injury was \$19,440 (including hospital, nursing home, emergency room, and home health care, but not doctors' services) (Rizzo et al. 1998).
- The total direct cost of all fall injuries for people 65 and older in 2000 was slightly more than \$19 billion: \$0.2 billion (\$179 million) for fatal falls, and \$19 billion for nonfatal falls (Stevens et al. 2006).
- By 2020, the annual direct and indirect cost of fall injuries is expected to reach \$43.8 billion (in current dollars) (Englander et al. 1996).

How do these costs break down?

Older Adult Falls

- [Preventing Falls Among Older Adults Brochures and Posters](#)
- [Figures and Maps](#)

Fact Sheets

- [Falls Among Older Adults: An Overview](#)
- [Costs of Falls Among Older Adults](#)
- [Hip Fractures Among Older Adults](#)
- [Falls in Nursing Homes](#)
- [CDC Fall Prevention Activities](#)

CenterCenter Overview **Age and sex**Organizational
ChartsInjury Center
LISTSERV

- The costs of fall injuries tend to increase with age (Stevens et al. 2006).
- In 2000, the costs of both fatal and nonfatal falls were higher for women than for men (Stevens et al. 2006).
- Medical costs for women, who made up 58% of older adults in 2000, were two to three times higher than for men (Stevens et al. 2006).

Contact CDC

Centers for Disease
Control and Prevention,
National Center for Injury
Prevention and Control
(NCIPC)
4770 Buford Hwy, NE
MS K-65
Atlanta, GA 30341-3717
Call:
1 (800) CDC-INFO (232-4636)
TTY: 1 (888) 232-6348
FAX: (770) 488-4760

Type of injury and treatment setting

- In 2000, nearly two thirds of the costs for nonfatal fall injuries were for those needing hospitalization. One fifth of costs were for injuries treated in emergency rooms (Stevens et al. 2006).
- Fractures were both the most common and most costly type of nonfatal injuries. Just over one third of nonfatal injuries were fractures, but they made up 61% of costs—or \$12 billion (Stevens et al. 2006).
- Hip fractures are the most frequent broken bones from falls. In the United States, 44% of direct health care costs for hip fractures are for hospitalization (Barrett-Connor 1995).
- Traumatic brain injuries and injuries to the hips, legs, and feet were the most common and costly fatal fall injuries in 2000. They made up 78% of fatalities and 79% of costs (Stevens et al. 2006).
- Injuries to internal organs caused 28% of deaths and accounted for 29% of costs from fatal falls (Stevens et al. 2006).

E-mail:
cdcinfo@cdc.gov

References

Alexander BH, Rivara FP, Wolf ME. The cost and frequency of hospitalization for fall-related injuries in older adults. *American Journal of Public Health* 1992;82(7):1020–3.

Barrett-Connor E. The economic and human costs of osteoporotic fracture. *American Journal of Medicine* 1995;98(suppl 2A):2A–3S to 2A–8S.

Englander F, Hodson TJ, Terregrossa RA. Economic dimensions of slip and fall injuries. *Journal of Forensic Science* 1996;41(5):733–46.

Hausdorff JM, Rios DA, Edelber HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Archives of Physical Medicine and Rehabilitation* 2001;82(8):1050–6.

Hornbrook MC, Stevens VJ, Wingfield DJ, Hollis JF, Greenlick MR, Ory MG. Preventing falls among community-dwelling older persons: results from a randomized trial. *The Gerontologist* 1994;34(1):16–23.

Rizzo JA, Friedkin R, Williams CS, Nabors J, Acampora D, Tinetti ME. Health care utilization and costs in a Medicare population by fall status. *Medical Care* 1998;36(8):1174–88.

Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and nonfatal falls among older adults. *Injury Prevention* 2006;12:290–5.

[Home](#) • [Policies and Regulations](#) • [Disclaimer](#) • [U.S. Government](#) • [FOIA](#) • [Contact Us](#)

Centers for Disease Control and Prevention, 1600 Clifton Rd., Atlanta, GA 30333, USA
Tel: 404-639-3311 • CDC Contact Center: 800-CDC-INFO • 888-232-6348 (TTY)

Department of Health
and Human Services
